


Lung Cancer Diagnosis for Primary Care

Daniel Nader, DO, FCCP


Cancer Treatment Center of America

Lung Cancer Facts

1 out of 14 people develop lung cancer.



Leading Sites of New Cancer Cases and Deaths: 2016 Estimates



Why are we here?

Prevention is the most effective strategy for reducing the burden of lung cancer

For non-preventable lung cancers,
Early detection & screening is critical

IMPROVE PATIENT CARE • SAVE LIVES



Sense of Urgency



**LUNG CANCER KILLS 433 PEOPLE
EVERY DAY!**

Patient Cohorts

- SYMPTOMATIC PATIENTS

- HIGH-RISK PATIENTS

- INCIDENTAL FINDINGS PATIENTS




Risk Factors

- Smoking
- Exposure to radon
- Exposure to other hazardous chemicals
 - Asbestos
 - Uranium
 - Arsenic
 - Cadmium
 - Chromium
 - Nickel
 - Some petroleum products
- Particle pollution
- Age
- Genetics: Personal or family history of lung cancer



Impact of Smoking



- Cigarette smoking is by far the most important risk factor for lung cancer
 - 80% of lung cancer deaths in women
 - 90% of lung cancer deaths in men
- Risk increases with both quantity and duration of smoking.
- Smoking
 - Cigarette smoking
 - Cigar & pipe smoking
 - Secondhand smoke
- Combined risk (*i.e.* smoking + asbestos)
- Smoking Cessation



Identifying Symptomatic Patients

- A cough that doesn't go away and gets worse over time
- A chronic cough or "smoker's cough"
- Hoarseness
- Constant chest pain
- Shortness of breath or wheezing
- Frequent lung infections such as bronchitis or pneumonia
- Coughing up blood
- Weight loss and loss of appetite
- Feeling tired or weak
- Infections such as bronchitis and pneumonia that don't go away or keep coming back



Identifying High-risk Patients

Two identifiers:

- Age 55-74


- 30-pack year Smoking History

Primary Care Provider Role

- Know the risk factors
- Counsel on smoking cessation
- Critical role in identifying:
 - Symptomatic patients
 - High-risk patients

Screening for Lung Cancer

National Lung Screening Trial (NLST)




20% fewer lung cancer deaths among 53,000 participants screened with low-dose helical (spiral) CT compared to those screened with chest X-rays.

The trial ended early due to promising results

NLST

Study Design




*High risk lung cancer was defined as a 30-year or more history of cigarette smoking. If the patient was a former smoker, they must have quit smoking within the last 15 years.

National Lung Screening Trial Research Team, et al. *N Engl J Med.* 2011;365:395-409.



NLST

Lung Cancer Mortality



- Positive screening rate for LDCT is 24.2%; for C X-RAY it is 6.9%
- 96.4% and 94.5% of the positive findings were false-positives in the LDCT and C X-RAY groups, respectively
- Lung cancer mortality was reduced by 20% in the LDCT group compared to the C X-RAY group
- All-cause mortality was reduced by 6.9% in the LDCT group compared with the C X-RAY group

National Lung Screening Trial Research Team, et al. *N Engl J Med.* 2011;365:395-409.



United States Preventive Task Force (USPSTF) Screening Recommendation

- Age 55-80
- ≥ 30 pack-year smoking history
- Current smoker or quit within the last 15 years
- Able and willing to receive treatment
- Screening should be discontinued
 - Once a person has not smoked for 15 years
 - Develops health problem(s) that substantially limits life expectancy or the ability or willingness to have curative lung surgery

Lung Screening Programs Requirements

Chart 6.1 Must-haves for screening sites

Follows an organized plan—a proven protocol—that is updated to include new technology and knowledge like that from NCCN

Has a high-quality screening program with enough staff and resources

Is accredited to do CT scans by a certifying organization, such as the American College of Radiology

Has scans read by an American Board of Radiology board-certified radiologist who's an expert in lung cancer screening

Has modern multislice CT equipment that does high-quality, low-dose, and non-contrast spiral CT

Is partnered with a health center that has: 1) experience and excellence in biopsy methods; 2) board-certified pulmonologists; and 3) board-certified thoracic surgeons who are experts in lung cancer





Case

- 58 year old male presented with dyspnea and dyspnea with exertion
- 40 pack year tobacco history, still smokes
- Father had lung lung cancer, age 68
- Hypertension
- ROS: chronic productive cough
- Exam: end expiratory wheezes

Case: Next Steps

- Treat exacerbation COPD
- PFT's
- Assess likelihood of lung cancer
- Manage pulmonary nodule



What are you going to tell the patient?

- A. You have a 5 mm nodule in the left lower lobe
- B. This might be cancer
- C. We want you to come back in 6 months
- D. We'll keep an eye on it
- E. A spot on the lung is always something to worry about. I don't want you to get frantic. We'll just take the test as we go along...



What's the patient thinking?

- What's a nodule? A “spot”. A “shadow”
- What's the chances that it is cancer?
- If it's not cancer, what else are we considering?
- Is it serious?
- What should I expect? Will I be able to breathe? Will I have pain?
- How long do we watch?



Tell the patient

- 1. Estimate the risk of cancer
- 2. Must be verbal. A letter is not satisfactory.
- 3. Avoid minimizing or dismissive language
- 4. Give this high priority, answer all the questions
- 5. Provide details of the nodule, it's the size of a pea, etc.
- 6. Explain the evaluation or surveillance
- 7. Let the patient know what to expect and acknowledge their concerns

What is a nodule?

- Solitary Pulmonary Nodule (SPN) is a radiographic opacity < 3 cm with at least 2/3 of its margins surrounded by lung parenchyma
- This excludes lymph nodes, atelectasis, and post-obstructive pneumonia
- Establishing the etiology of a SPN assumes critical importance



SPN

- 150,00 per year
- CT imaging: 8% to 51%
- Prevalence of malignancy: 1.1% to 12%
- Accurate and timely diagnosis is important as treatment of early stage lung cancer provides the highest chance for cure.



Clinical Evaluation: History

- History of smoking
- History of prior malignancy
- History of interstitial Lung Disease
- Residence in or travel to areas endemic with fungal pathogens
- Prior CT scan or Chest radiograph



CT Scan

- Thin sections
- 1.5 mm
- Lung and mediastinal windows
- Contrast for mediastinal structures and nodule enhancement



Growth Rate

- Malignant nodules double in 20 to 400 days
- Volume of a sphere = $4\pi r^3/3$
- An increase in diameter of 26% represents a doubling in volume
- Example: 4mm nodule to 5mm is almost a doubling in volume
- A solid nodule which does not change in size over 2 years is considered benign



Size Matters

- Size trumps morphology: diameter and likelihood of cancer in screened smokers


- < 3 mm: 0.2%
- 4 – 7 mm 2.7%
- 8 – 30 mm 18%
- > 30 mm 99%



Benign Patterns of Calcifications



diffuse



central




popcorn




concentric


Indeterminate Patterns of Calcifications



stippled



eccentric




**inhomogeneous
calcification
of irregular nodule**

Findings that suggest a benign lesion

- Shape: oblong, polygonal, concave margins
- Edge smooth
- Subpleural location (attached to the pleura)
- Satellite nodules

Oblong, polygonal, concave margins




follow-up

Ground Glass Opacities



- Subsolid nodule = pure GGO
- Partly solid GGN: focal nodular area of increased attenuation
- Semi-solid GGN: may represent a histologic spectrum of adenocarcinoma
- Atypical adenomatous hyperplasia → adenocarcinoma in situ → minimally invasive adenocarcinoma → lepidic predominant adenocarcinoma

Ground-glass opacity



5 mm, standard algorithm

Ground-glass opacity Bronchioloalveolar Carcinoma



Lepidic growth

Which nodule is malignant?





PET Scan

- Combined PET-CT correlates results from 2 modalities
- Glucose analog tagged with positron-emitting isotope fluorine(FDG)
- Reveals aspects of tumor function and metabolism
- Metabolic activity quantitated using Standard Uptake Value (SUV) Mean value > 2.5



Case 2: PET



Pulmonary nodules: FDG-PET diagnosis

- Meta-analysis of 40 studies with 1474 nodules
- SUV 2.5
- Sensitivity 97%; specificity 78%; accuracy 90%

Pulmonary nodules: FDG-PET diagnosis

- 625 patients
- 433 diagnosed with cancer
- False negatives:
 - 11/23 (48%) Lepidic predominant adenocarcinoma
 - 8/234 (3%) adenocarcinoma
 - 4/14 (29%) carcinoid
 - 2/8 (25%) renal cell metastases

Pulmonary nodules: FDG-PET diagnosis

- 136 nodules < 3 cm; 81 malignant
- All 20 < 1 cm (8 malignant) were negative on PET
- 101 solid nodules 1-3 cm (63 malignant) sensitivity 90%, specificity 71%
- 15 ground-glass nodules 1-3 cm (10 malignant) sensitivity 10%, specificity 20%

Solitary Pulmonary Nodule Calculator

http://www.chestx-ray.com/SPN/SPNProb.html

Probability of Malignancy in SPN: Logistic Regression

Compare the above result to this method derived from multivariate logistic regression in 629 patients (65% benign, 23% malignant, 12% indeterminate). The equation is based on 3 clinical and 3 radiographic variables.

$$\text{Probability of Malignancy} = e^x / (1 + e^x)$$


where $x = -6.8272 + (0.0391 * \text{Age}) + (0.7917 * \text{Cigarettes}) + (1.3388 * \text{Cancer}) + (0.1274 * \text{Diameter}) + (1.0407 * \text{Spiculation}) + (0.7838 * \text{Upper})$.

Note: this equation is not applicable to patients with a diagnosis of cancer that has been made within the previous 5 years or to patients with previous lung cancer.

Reference:

Swensen SJ, Silverstein MD, Ilstrup DM, Schleck CD, Edell ES. [The probability of malignancy in solitary pulmonary nodules. Application to small radiologically indeterminate nodules.](#) Arch Intern Med 1997; 157:849-855.
[\[Related Records\]](#)

Age (yrs):	<input type="text" value="62"/>
SPN diameter (mm):	<input type="text" value="25"/>
Edge:	<input type="text" value="Spiculated"/>
Previous Malignancy:	<input type="text" value="No Hx Malignancy"/>
Smoking:	<input type="text" value="Smoker or Former Smoker"/>
Location:	<input type="text" value="Upper lobe"/>
<input type="button" value="Calculate Probability of Malignancy"/>	
The Probability of Malignancy is:	<input type="text" value="80"/>
<input type="button" value="Reset"/>	



Probability of Cancer in Pulmonary Nodules

- Predictors of cancer model
 - Older age
 - Female sex
 - Family history of lung cancer
 - Emphysema
 - Larger nodule size
 - Location of nodule in the upper lobe
 - Part-solid nodule type
 - Lower nodule count
 - Spiculation

Probability of Cancer in Pulmonary Nodules

Input:

Age years

Sex Female (0.6011)
 Male (0)

Family history of lung cancer (0.2961)

Emphysema (0.2953)

Nodule size mm

Nodule type Nonsolid or ground-glass (-0.1276)
 Partially solid (0.377)
 Solid (0)

Nodule in upper lung (0.6581)

Nodule count #

Spiculation (0.7729)

Results:

Log odds

Cancer probability %

Decimal precision



Algorithmic Approach to SPN

- Review previous imaging
- Solid or subsolid
- Solid nodules < 8 mm can be followed
- Solid nodules > 8 mm need pretest probability for cancer
- Pretest probability < 5%, follow
- Pretest probability > 60% needs tissue diagnosis
- Intermediate range: PET



Where Do You Compare With Your Peers?

- Annals of the American Thoracic Society, January 2018
- An Assessment of Primary Care and Pulmonary Provider Perspectives on Lung Cancer Screening

Primary Provider Perspective On Lung Cancer Screening

- 196 participants
- 80% primary care
- 41% University or affiliated clinics
- 47% county hospital based clinics




- 74% endorsed screening effectiveness
- Key barriers: inadequate time (36%), inadequate staffing (36%), patients have too many other illnesses to address (38%)
- Decision aids were important, at point of referral, to facilitate screening
- Point-of-care referral materials may be helpful in reducing knowledge gaps and clinical burden of referral

Lung Cancer Screening Recommendation


RISK CRITERIA	USPSTF	CMS	NCCN
Age	55-80	55-77	55
Pack-year Smoking History	≥ 30	≥ 30	≥ 30
Status	Current smoker or quit within the last 15 years	Current smoker or quit within the last 15 years	Current smoker or quit within the last 15 years
Other	Able and willing to receive treatment	Shared decision making	Shared decision making
Discontinuation Guidelines	<p>Age 80</p> <p>Once a person has not smoked for 15 years</p> <p>Develops health problem(s) that substantially limits life expectancy or the ability or willingness to have curative lung surgery</p>	Age 77	N/A

Bronchoscopy Reach Limitations

Flexible
Bronchoscopy



Reach of Current Day
Bronchoscopic Vision &
Control





Overview




How Might Robotics Help?

- Not restricted to conventional scope design
 - Possible to reinvent the scope?
- Incorporate latest guided approaches
- Improved instrumentation





Scope Tip



Bronchoscopy Today





My Patient





Genomic Test Results





Questions?

